

## General

### Title

Stroke: percentage of ischemic stroke patients who develop a symptomatic intracranial hemorrhage within (less than or equal to) 36 hours after the onset of treatment with IV or IA thrombolytic (t-PA) therapy, or mechanical endovascular reperfusion procedure.

### Source(s)

The Joint Commission. Disease-specific care certification program. Comprehensive stroke: performance measurement implementation guide. Oakbrook Terrace (IL): The Joint Commission; 2015 Mar. 278 p.

## Measure Domain

### Primary Measure Domain

Clinical Quality Measures: Outcome

### Secondary Measure Domain

Does not apply to this measure

## Brief Abstract

### Description

This measure is used to assess the percentage of ischemic stroke patients who develop a symptomatic intracranial hemorrhage (i.e., clinical deterioration greater than or equal to 4 point increase on National Institutes of Health Stroke Scale [NIHSS] and brain image finding of parenchymal hematoma, or subarachnoid hemorrhage, or intraventricular hemorrhage) within (less than or equal to) 36 hours after the onset of treatment with intravenous (IV) or intra-arterial (IA) thrombolytic (t-PA) therapy, or mechanical endovascular reperfusion procedure (i.e., mechanical endovascular thrombectomy with a clot retrieval device).

This measure represents the overall rate. The following rates are also reported:

Hemorrhagic transformation for patients treated with intra-venous (IV) thrombolytic (t-PA) therapy only

Hemorrhagic transformation for patients treated with intra-arterial (IA) thrombolytic (t-PA) therapy or mechanical endovascular reperfusion therapy

## Rationale

Intravenous (IV) thrombolytic (t-PA) therapy for acute ischemic stroke was approved by the U.S. Food and Drug Administration in 1996, following findings from the National Institute of Neurological Disorders and Stroke (NINDS) trial which demonstrated favorable outcomes in 31% to 50% of patients treated with recombinant tissue plasminogen activator (r-tPA), as compared to 20% to 38% of patients treated with placebo. Intra-arterial (IA) t-PA therapy has since been used to improve recanalization and clinical outcomes for select patients nonresponsive to IV therapy. Intracranial hemorrhage is the major risk of t-PA therapy with similar rates reported for both IV and IA routes. The NINDS trial found that 6.4% of patients treated with IV t-PA experienced symptomatic bleeding. Findings from the Prolyse in Acute Cerebral Thromboembolism (PROACT II) study found the intracranial hemorrhage with neurological deterioration within 24 hours occurred in 10% of patients treated with IA recombinant prourokinase. In addition to these agents, other available thrombolytic drugs include: streptokinase, p-anisoylated lys-plasminogen-streptokinase activator, and urokinase.

Endovascular reperfusion therapy in acute ischemic stroke comprises a number of pharmacological and mechanical procedures. Mechanical endovascular thrombectomy is a treatment option for patients with large vessel occlusions in whom pharmacological thrombolysis is contraindicated or might be ineffective. A number of mechanical endovascular thrombectomy devices, also known as clot retrieval devices, are currently undergoing clinical evaluation. Mechanical endovascular thrombectomy devices are intended to improve tissue rescue and diminish reperfusion hemorrhage while broadening the population eligible for therapy. These devices may be used alone or in conjunction with chemical thrombolysis (i.e., IV or IA t-PA).

## Evidence for Rationale

Adams HP Jr, Brott TG, Furlan AJ, Gomez CR, Grotta J, Helgason CM, Kwiatkowski T, Lyden PD, Marler JR, Torner J, Feinberg W, Mayberg M, Thies W. Guidelines for thrombolytic therapy for acute stroke: a supplement to the guidelines for the management of patients with acute ischemic stroke. A statement for healthcare professionals from a Special Writing Group of the Stroke Council. *Circulation*. 1996 Sep 1;94(5):1167-74. [102 references] [PubMed](#)

Adams HP, del Zoppo G, Alberts MJ, Bhatt DL, Brass L, Furlan A, Grubb RL, Higashida RT, Jauch EC, Kidwell C, Lyden PD, Morgenstern LB, Qureshi AI, Rosenwasser RH, Scott PA, Wijdicks EF, American Heart Association, American Stroke Association Stroke Council, Clinical Cardiology Council, Cardiovascular Radiology and Intervention Council, Atherosclerotic Peripheral Vascular Disease and Quality of Care Outcomes in Research Interd[TRUNC]. Guidelines for the early management of adults with ischemic stroke: a guideline from the American Heart Association/American Stroke Association Stroke Council, Clinical Cardiology Council, Cardiovascular Radiology and Intervention Council, and the Atherosclerotic Peripheral Vascular Disease and Quality of Care Outcomes in Research Interdisciplinary Working Groups: the American Academy of Neurology affirms the value of this guideline as an educational tool for neurologists. *Stroke*. 2007 May;38(5):1655-711. [PubMed](#)

Broderick JP, Palesch YY, Demchuk AM, Yeatts SD, Khatri P, Hill MD, Jauch EC, Jovin TG, Yan B, Silver FL, von Kummer R, Molina CA, Demaerschalk BM, Budzik R, Clark WM, Zaidat OO, Malisch TW, Goyal M, Schonewille WJ, Mazighi M, Engelter ST, Anderson C, Spilker J, Carrozzella J, Ryckborst KJ, Janis LS, Martin RH, Foster LD, Tomsick TA, Interventional Management of Stroke (IMS) III Investigators. Endovascular therapy after intravenous t-PA versus t-PA alone for stroke. *N Engl J Med*. 2013 Mar 7;368(10):893-903. [PubMed](#)

Ciccone A, Valvassori L, Nichelatti M, Sgoifo A, Ponzio M, Sterzi R, Boccardi E, SYNTHESIS Expansion Investigators. Endovascular treatment for acute ischemic stroke. *N Engl J Med*. 2013 Mar 7;368(10):904-13. [PubMed](#)

del Zoppo GJ, Higashida RT, Furlan AJ, Pessin MS, Gent M, Driscoll RM, PROACT Investigators. The Prolyse in Acute Cerebral Thromboembolism Trial (PROACT): results of 6 mg dose tier. *Stroke*. 1996;27:164.

del Zoppo GJ, Higashida RT, Furlan AJ. The case for a phase III trial of cerebral intraarterial fibrinolysis. *AJNR Am J Neuroradiol*. 1994;15:1217-22.

Donnan GA, Davis SM, Chambers BR, Gates PC, Hankey GJ, McNeil JJ, Rosen D, Stewart-Wynne EG, Tuck RR. Trials of streptokinase in severe acute ischaemic stroke. *Lancet*. 1995 Mar 4;345(8949):578-9. [PubMed](#)

Fibrinolytic Therapy Trialists' (FFT) Collaborative Group. Indications for fibrinolytic therapy in suspected acute myocardial infarction: collaborative overview of early mortality and major morbidity results from all randomised trials of more than 1000 patients. *Lancet*. 1994 Feb 5;343(8893):311-22. [58 references] [PubMed](#)

Hacke W, Kaste M, Fieschi C, Toni D, Lesaffre E, von Kummer R, Boysen G, Bluhmki E, Hoxter G, Mahagne MH, et al. Intravenous thrombolysis with recombinant tissue plasminogen activator for acute hemispheric stroke. The European Cooperative Acute Stroke Study (ECASS). *JAMA*. 1995 Oct 4;274(13):1017-25. [PubMed](#)

Jauch EC, Saver JL, Adams HP, Bruno A, Connors JJ, Demaerschalk BM, Khatri P, McMullan PW, Qureshi AI, Rosenfield K, Scott PA, Summers DR, Wang DZ, Wintermark M, Yonas H, American Heart Association Stroke Council, Council on Cardiovascular Nursing, Council on Peripheral Vascular Disease, Council on Clinical Cardiology. Guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2013 Mar;44(3):870-947. [PubMed](#)

Kidwell CS, Jahan R, Gornbein J, Alger JR, Nenov V, Ajani Z, Feng L, Meyer BC, Olson S, Schwamm LH, Yoo AJ, Marshall RS, Meyers PM, Yavagal DR, Wintermark M, Guzy J, Starkman S, Saver JL, MR RESCUE Investigators. A trial of imaging selection and endovascular treatment for ischemic stroke. *N Engl J Med*. 2013 Mar 7;368(10):914-23. [PubMed](#)

Koh JS, Lee SJ, Ryu CW, Kim HS. Safety and efficacy of mechanical thrombectomy with solitaire stent retrieval for acute ischemic stroke: a systematic review. *Neurointervention*. 2012 Feb;7(1):1-9. [PubMed](#)

Leifer D, Bravata DM, Connors JJ, Hinchey JA, Jauch EC, Johnston SC, Latchaw R, Likosky W, Ogilvy C, Qureshi AI, Summers D, Sung GY, Williams LS, Zorowitz R, American Heart Association Special Writing Group of the Stroke Council, Atherosclerotic Peripheral Vascular Disease Working Group, Council on Cardiovascular Surgery and Anesthesia, Council on Cardiovascular Nursing. Metrics for measuring quality of care in comprehensive stroke centers: detailed follow-up to Brain Attack Coalition comprehensive stroke center recommendations: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2011 Mar;42(3):849-77. [PubMed](#)

Levy DE, Brott TG, Haley EC, Marler JR, Sheppard GL, Barsan W, Broderick JP. Factors related to intracranial hematoma formation in patients receiving tissue-type plasminogen activator for acute ischemic stroke. *Stroke*. 1994 Feb;25(2):291-7.

Marder VJ, Sherry S. Thrombolytic therapy: current status (1). *N Engl J Med*. 1988 Jun 9;318(23):1512-20. [PubMed](#)

Menon BK, Saver JL, Prabhakaran S, Reeves M, Liang L, Olson DM, Peterson ED, Hernandez AF, Fonarow GC, Schwamm LH, Smith EE. Risk score for intracranial hemorrhage in patients with acute ischemic stroke treated with intravenous tissue-type plasminogen activator. *Stroke*. 2012 Sep;43(9):2293-9. [PubMed](#)

Nogueira RG, Lutsep HL, Gupta R, Jovin TG, Albers GW, Walker GA, Liebeskind DS, Smith WS, TREVO 2 Trialists. Trevo versus Merci retrievers for thrombectomy revascularisation of large vessel occlusions in acute ischaemic stroke (TREVO 2): a randomised trial. *Lancet*. 2012 Oct 6;380(9849):1231-40. [PubMed](#)

Randomised controlled trial of streptokinase, aspirin, and combination of both in treatment of acute ischaemic stroke. Multicentre Acute Stroke Trial--Italy (MAST-I) Group. Lancet. 1995 Dec 9;346(8989):1509-14. [PubMed](#)

Rubiera M, Ribo M, Pagola J, Coscojuela P, Rodriguez-Luna D, Maisterra O, Ibarra B, Piñeiro S, Meler P, Romero FJ, Alvarez-Sabin J, Molina CA. Bridging intravenous-intra-arterial rescue strategy increases recanalization and the likelihood of a good outcome in nonresponder intravenous tissue plasminogen activator-treated patients: a case-control study. Stroke. 2011 Apr;42(4):993-7. [PubMed](#)

Saver JL, Jahan R, Levy EI, Jovin TG, Baxter B, Nogueira RG, Clark W, Budzik R, Zaidat OO, SWIFT Trialists. Solitaire flow restoration device versus the Merci Retriever in patients with acute ischaemic stroke (SWIFT): a randomised, parallel-group, non-inferiority trial. Lancet. 2012 Oct 6;380(9849):1241-9. [PubMed](#)

Sims JR, Gharai LR, Schaefer PW, Vangel M, Rosenthal ES, Lev MH, Schwamm LH. ABC/2 for rapid clinical estimate of infarct, perfusion, and mismatch volumes. Neurology. 2009 Jun 16;72(24):2104-10. [PubMed](#)

Sloan MA, Price TR, Petito CK, Randall AM, Solomon RE, Terrin ML, Gore J, Collen D, Kleiman N, Feit F, et al. Clinical features and pathogenesis of intracerebral hemorrhage after rt-PA and heparin therapy for acute myocardial infarction: the Thrombolysis in Myocardial Infarction (TIMI) II Pilot and Randomized Clinical Trial combined experience. Neurology. 1995 Apr;45(4):649-58. [PubMed](#)

Smith WS, Sung G, Saver J, Budzik R, Duckwiler G, Liebeskind DS, Lutsep HL, Rymer MM, Higashida RT, Starkman S, Gobin YP, Multi MERCI Investigators, Frei D, Grobelny T, Hellinger F, Huddle D, Kidwell C, Koroshetz W, Marks M, Nesbit G, Silverman IE. Mechanical thrombectomy for acute ischemic stroke: final results of the Multi MERCI trial. Stroke. 2008 Apr;39(4):1205-12. [PubMed](#)

The Joint Commission. Disease-specific care certification program. Comprehensive stroke: performance measurement implementation guide. Oakbrook Terrace (IL): The Joint Commission; 2015 Mar. 278 p.

Tissue plasminogen activator for acute ischemic stroke. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. N Engl J Med. 1995 Dec 14;333(24):1581-7. [PubMed](#)

## Primary Health Components

Ischemic stroke; intracranial hemorrhage; intra-venous (IV) thrombolytic (t-PA) therapy; intra-arterial (IA) t-PA therapy; mechanical endovascular reperfusion therapy

## Denominator Description

Ischemic stroke patients treated with intravenous (IV) thrombolytic (t-PA) therapy only (IVO) or intra-arterial (IA) t-PA therapy, or who undergo mechanical endovascular reperfusion therapy (see the related "Denominator Inclusions/Exclusions" field)

## Numerator Description

Ischemic stroke patients who develop a symptomatic intracranial hemorrhage less than or equal to 36 hours after the onset of treatment with intravenous (IV) thrombolytic (t-PA) therapy, or intra-arterial (IA) t-PA therapy, or mechanical endovascular reperfusion therapy

## Evidence Supporting the Measure

## Type of Evidence Supporting the Criterion of Quality for the Measure

A clinical practice guideline or other peer-reviewed synthesis of the clinical research evidence

A formal consensus procedure, involving experts in relevant clinical, methodological, public health and organizational sciences

A systematic review of the clinical research literature (e.g., Cochrane Review)

One or more research studies published in a National Library of Medicine (NLM) indexed, peer-reviewed journal

## Additional Information Supporting Need for the Measure

Unspecified

## Extent of Measure Testing

Unspecified

## State of Use of the Measure

### State of Use

Current routine use

### Current Use

not defined yet

## Application of the Measure in its Current Use

### Measurement Setting

Hospital Inpatient

### Professionals Involved in Delivery of Health Services

not defined yet

### Least Aggregated Level of Services Delivery Addressed

Single Health Care Delivery or Public Health Organizations

### Statement of Acceptable Minimum Sample Size

Specified

## Target Population Age

Age greater than or equal to 18 years

## Target Population Gender

Either male or female

# National Strategy for Quality Improvement in Health Care

## National Quality Strategy Aim

Better Care

## National Quality Strategy Priority

Making Care Safer

Prevention and Treatment of Leading Causes of Mortality

# Institute of Medicine (IOM) National Health Care Quality Report Categories

## IOM Care Need

Getting Better

## IOM Domain

Effectiveness

Safety

# Data Collection for the Measure

## Case Finding Period

Unspecified

## Denominator Sampling Frame

Patients associated with provider

## Denominator (Index) Event or Characteristic

Clinical Condition

Institutionalization

Patient/Individual (Consumer) Characteristic

Therapeutic Intervention

## Denominator Time Window

not defined yet

## Denominator Inclusions/Exclusions

### Inclusions

Discharges with International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) Principal Diagnosis Code for ischemic stroke as defined in the appendices of the original measure documentation

AND

Patients with documented thrombolytic (intravenous [IV] or intra-arterial [IA] thrombolytic [t-PA]) therapy (ICD-9-CM Principal or Other Procedure Codes as defined in the appendices of the original measure documentation)

OR

Patients with documented mechanical endovascular reperfusion therapy (ICD-9 CM Principal or Other Procedure Codes as defined in the appendices of the original measure documentation)

### Exclusions

Patients less than 18 years of age

Patients who have a Length of Stay greater than 120 days

Patients admitted for *Elective Carotid Intervention* (as defined in the Data Elements)

Patients transferred to the hospital following treatment with IV t-PA therapy or IA t-PA therapy or mechanical endovascular reperfusion therapy initiated prior to arrival at the hospital

Patients who hemorrhage prior to the onset of treatment with IV t-PA therapy or IA t-PA therapy or mechanical endovascular reperfusion therapy

## Exclusions/Exceptions

not defined yet

## Numerator Inclusions/Exclusions

### Inclusions

Ischemic stroke patients who develop a symptomatic intracranial hemorrhage less than or equal to 36 hours after the onset of treatment with intravenous (IV) thrombolytic (t-PA) therapy, or intra-arterial (IA) t-PA therapy, or mechanical endovascular reperfusion therapy

### Exclusions

None

## Numerator Search Strategy

Institutionalization

## Data Source

Administrative clinical data

Paper medical record

## Type of Health State

Adverse Health State

## Instruments Used and/or Associated with the Measure

- The National Institutes of Health Stroke Scale (NIHSS)
- Comprehensive Stroke (CSTK) Initial Patient Population Algorithm Flowchart
- CSTK-05: Hemorrhagic Transformation (Overall Rate) Flowchart

## Computation of the Measure

### Measure Specifies Disaggregation

Measure is disaggregated into categories based on different definitions of the denominator and/or numerator

### Basis for Disaggregation

The CSTK-05 measure is reported as an overall rate which includes ischemic stroke patients who develop a symptomatic hemorrhage after reperfusion therapy.

CSTK-05a: Ischemic stroke patients who develop a symptomatic intracranial hemorrhage less than or equal to 36 hours after the onset of treatment with IV thrombolytic (t-PA) therapy only (IVO).

CSTK-05b: Ischemic stroke patients who develop a symptomatic intracranial hemorrhage less than or equal to 36 hours after the onset of treatment with IA thrombolytic (t-PA) therapy or mechanical endovascular reperfusion therapy.

CSTK-05a and CSTK-05b are subsets of the overall rate, and stratified by the type of therapy.

Data Reported As: Aggregate rate generated from count data reported as a proportion.

## Scoring

Rate/Proportion

## Interpretation of Score

Desired value is a lower score

## Allowance for Patient or Population Factors



not defined yet

## Description of Allowance for Patient or Population Factors

Risk adjustment for this measure is applied to the following data elements:

Admission Date  
Birthdate  
Hispanic Ethnicity  
International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) Other  
Diagnosis Codes  
Intravenous (IV) Thrombolytic Therapy Prior to Intra-arterial (IA) or Mechanical Reperfusion Therapy  
Initial Blood Glucose Value at Hospital Arrival  
Initial Blood Pressure at Hospital Arrival  
Initial National Institutes of Health Stroke Scale (NIHSS) Score at Hospital Arrival  
Initial Platelet Count at Hospital Arrival  
Race  
Sex

## Standard of Comparison

not defined yet

## Identifying Information

### Original Title

CSTK-05: hemorrhagic transformation (overall rate).

### Measure Collection Name

Advanced Certification in Disease-specific Care Measures

### Measure Set Name

Comprehensive Stroke Standardized Performance Measures

### Submitter

The Joint Commission - Health Care Accreditation Organization

### Developer

The Joint Commission - Health Care Accreditation Organization

### Funding Source(s)

All external funding for measure development has been received and used in full compliance with The Joint Commission's corporate sponsorship policies, which are available upon written request to The Joint

Commission.

## Composition of the Group that Developed the Measure

Unspecified

## Financial Disclosures/Other Potential Conflicts of Interest

Expert panel members have made full disclosure of relevant financial and conflict of interest information in accordance with The Joint Commission's conflict of interest policies, copies of which are available upon written request The Joint Commission.

## Adaptation

This measure was not adapted from another source.

## Date of Most Current Version in NQMC

2015 Mar

## Measure Maintenance

This measure is reviewed and updated by the developing organization every 6 months.

## Date of Next Anticipated Revision

2015 Jul

## Measure Status

This is the current release of the measure.

The measure developer reaffirmed the currency of this measure in April 2016.

## Measure Availability

Source available from [The Joint Commission Web site](#) .

For more information, contact The Joint Commission at One Renaissance Blvd., Oakbrook Terrace, IL 60181; Phone: 630-792-5800; Fax: 630-792-5005; Web site: [www.jointcommission.org](http://www.jointcommission.org)

.

## NQMC Status

This NQMC summary was completed by ECRI Institute on May 19, 2015. The information was verified by the measure developer on June 22, 2015.

The information was reaffirmed by the measure developer on April 6, 2016.

## Copyright Statement

This NQMC summary is based on the original measure, which is subject to the measure developer's copyright restrictions.

## Production

### Source(s)

The Joint Commission. Disease-specific care certification program. Comprehensive stroke: performance measurement implementation guide. Oakbrook Terrace (IL): The Joint Commission; 2015 Mar. 278 p.

## Disclaimer

### NQMC Disclaimer

The National Quality Measures Clearinghouse<sup>®</sup> (NQMC) does not develop, produce, approve, or endorse the measures represented on this site.

All measures summarized by NQMC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public and private organizations, other government agencies, health care organizations or plans, individuals, and similar entities.

Measures represented on the NQMC Web site are submitted by measure developers, and are screened solely to determine that they meet the [NQMC Inclusion Criteria](#).

NQMC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or its reliability and/or validity of the quality measures and related materials represented on this site. Moreover, the views and opinions of developers or authors of measures represented on this site do not necessarily state or reflect those of NQMC, AHRQ, or its contractor, ECRI Institute, and inclusion or hosting of measures in NQMC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding measure content are directed to contact the measure developer.